



Filler Dissolving Post Treatment Advice

Hyalase is an enzyme which breaks down hyaluronic acid fillers, but it can also break down naturally occurring hyaluronic acid present in the body. The results can be unpredictable and the effect dramatic with possible loss of volume and some skin laxity. Although some of the effects can be immediate, it can take up to 2 weeks for the final results to be seen and the procedure may need to be repeated.

Hyalase administration can result in anaphylaxis (a severe allergic reaction) which in itself is life threatening and requires immediate medical attention. Symptoms of a severe allergic reaction can include shortness of breath, wheezing, coughing, difficulty swallowing, swelling of the tongue, eyelids, lips, hoarseness of the voice, stomach pain, nausea or diarrhoea.

If you have any of the above symptoms please report to your nearest Accident and Emergency Department or call 999 for an ambulance.

After treatment

After the procedure some other common injection-related reactions might occur. These reactions include redness, swelling, pain, itching, bruising and tenderness at the injection site. They have generally been described as mild to moderate and typically resolve spontaneously after a few days after the injection. Bruising may occasionally be more significant.

If you have any concerns following treatment, do not hesitate to contact the clinic as soon as possible.

Should you experience any unexpected side effects or any that concern you, please contact the clinic as soon as possible.

Please ensure you discuss your overall treatment programme with your clinician as this will enable you to receive the best results from treatment.

Est-Ethics
170 Ingram St
Italian Centre
Glasgow
G11DN

0141 343 24 24
admin@est-ethics.com

www.est-ethics.com

